

Two-dimensional gravity with an invariant energy scale and arbitrary dilaton potential

S. Mignemi*

Dipartimento di Matematica, Università di Cagliari,
Via Ospedale 72, 09124 Cagliari, Italy

INFN, Sezione di Cagliari

Abstract

We investigate a model of two-dimensional gravity with arbitrary scalar potential obtained by gauging a deformation of de Sitter or more general algebras, which accounts for the existence of an invariant energy scale. We obtain explicit solutions of the field equations and discuss their properties.

*email: smignemi@unica.it

Deformations of the Poincaré algebra allowing for the existence of a fundamental length (energy) scale have recently been the subject of several investigations [1, 2]. Their interest is motivated by the fact that theories of quantum gravity seem to imply the existence of a minimal length scale of the order of the Planck length [3].

One may wonder whether the framework of deformed Poincaré algebra can be extended to include gravitational interactions, giving rise in this way to an effective modification of general relativity at small scales. As is well known, a theory of gravity can be obtained in general by gauging the space-time symmetry group, and this approach was adopted in [4] for constructing a two-dimensional gravitational model invariant under the specific deformation of Poincaré algebra introduced in ref. [2].

In this letter we go a step further, extending the model of [4] to the case where a dilaton potential is present [5]. This class of two-dimensional theories includes between others (anti-)de Sitter [6], or extended Poincaré models [7]. Finding suitable deformations of the algebras of [5] compatible with the assumptions of [2] is not trivial, because the Jacobi identities impose constraints on the form of the deformed algebra.

The deformed Poincaré algebra of [2] is given in two dimensions by the commutation relations

$$[P_a, P_b] = 0, \quad [J, P_0] = P_1 - \frac{P_0 P_1}{\kappa}, \quad [J, P_1] = P_0 - \frac{P_1^2}{\kappa}, \quad (1)$$

where P_a are the generators of translations and J that of boosts and $a = 0, 1$. Tangent space indices are lowered and raised by the tensor $h_{ab} = \text{diag}(-1, 1)$. We also make use of the antisymmetric tensor ϵ_{ab} , with $\epsilon_{01} = 1$. The deformation parameter κ has the dimension of a mass and can be identified with the inverse of the Planck length. The algebra (1) can be considered as an example of nonlinear algebra [8].

A gravitational model which is invariant under the local symmetry (1) was studied in [4] and is described by a lagrangian

$$L = \eta_2 R + \eta_a T^a + \frac{\eta_1}{\kappa} \eta_a \omega \wedge e^a, \quad (2)$$

where $R = d\omega$ is the curvature and $T^a = de^a + \epsilon^a_b \omega \wedge e^b$ the torsion and η_a , η_2 are a triplet of scalar fields in the coadjoint representation of the algebra. η_2 is often called dilaton.

We want to generalize this two-dimensional model to include dilaton-dependent potentials $\Lambda(\eta_2)$, analogous to those studied in [5] in the Lorentz-invariant case. Simple special cases were de Sitter gravity [6], corresponding to $\Lambda = \lambda\eta_2$, or extended Poincaré gravity [7], corresponding to $\Lambda = \lambda$. In these models, the commutation relation between the generators of translations P_a is of the form [5]

$$[P_0, P_1] = \Lambda(J), \quad (3)$$

where Λ is the same function as the potential, and J is the generator of boosts.

Unfortunately, it is not possible to simply change the momentum commutation relation in (1) to the form (3) because the set does not satisfy the generalized Jacobi identities of nonlinear algebras [8]. The simplest generalization of the algebra (1) is to admit a relation of the form $[P_0, P_1] = f(P_a)\Lambda(J)$. A simple calculation shows that the Jacobi identities are in fact satisfied if $f(P_a) = \left(1 - \frac{P_0}{\kappa}\right)^3$. Hence, we shall consider the nonlinear algebra

$$[P_a, P_b] = \left(1 - \frac{P_0}{\kappa}\right)^3 \Lambda(J), \quad [J, P_0] = P_1 - \frac{P_0 P_1}{\kappa}, \quad [J, P_1] = P_0 - \frac{P_1^2}{\kappa}. \quad (4)$$

For $\kappa \rightarrow \infty$, the algebra reduces to that obtained in [5] for the Lorentz-invariant limit. The algebra (4) admits a Casimir invariant

$$C = \frac{P_1^2 - P_0^2}{\left(1 - \frac{P_0}{\kappa}\right)^2} + 2 \int \Lambda(J) dJ. \quad (5)$$

In order to construct a gauge theory for this algebra, we proceed as in [4] and adopt the formalism of Ikeda [8]. Given an algebra with commutation relations $[T_A, T_B] = W_{AB}(T)$, one introduces gauge fields A^A and a coadjoint multiplet of scalar fields η_A , which under infinitesimal transformations of parameter ξ^A transform as

$$\begin{aligned} \delta A^A &= d\xi^A + U_{BC}^A(\eta) A^B \xi^C, \\ \delta \eta_A &= -W_{AB}(\eta) \xi^B, \end{aligned} \quad (6)$$

where U_{BC}^A and W_{AB} are functions of the fields η_A which satisfy

$$U_{BC}^A = \frac{\partial W_{BC}}{\partial \eta_A}. \quad (7)$$

In our case $A = 0, 1, 2$, and we identify T^a with P^a and T^2 with J . In analogy with [6, 7], we can then also identify A^a with the zweibeins e^a and A^2 with the spin connection ω , and obtain a gauge theory for gravity in two dimensions. A gauge-invariant lagrangian can then be obtained with the methods of [8], and reads

$$L = \eta_2 R + \eta_a T^a + \frac{1}{2} \left(1 - \frac{\eta_0}{\kappa}\right)^3 \Lambda(\eta_2) \epsilon_{ab} e^a \wedge e^b + \frac{\eta_1}{\kappa} \eta_a \omega \wedge e^a, \quad (8)$$

with field equations

$$\begin{aligned} d\eta_0 &= \left(\eta_1 - \frac{\eta_1 \eta_0}{\kappa}\right) \omega - \left(1 - \frac{\eta_0}{\kappa}\right)^3 \Lambda(\eta_2) e^1, \\ d\eta_1 &= \left(\eta_0 - \frac{\eta_1^2}{\kappa}\right) \omega + \left(1 - \frac{\eta_0}{\kappa}\right)^3 \Lambda(\eta_2) e^0, \\ d\eta_2 &= -\left(\eta_1 - \frac{\eta_1 \eta_0}{\kappa}\right) e^0 - \left(\eta_0 - \frac{\eta_1^2}{\kappa}\right) e^1, \\ T^0 &= \omega \wedge \frac{\eta_1}{\kappa} e^0 + \frac{3}{\kappa} \left(1 - \frac{\eta_0}{\kappa}\right)^2 \Lambda(\eta_2) e^0 \wedge e^1, \\ T^1 &= \omega \wedge \left(\frac{\eta_0}{\kappa} e^0 + \frac{2\eta_1}{\kappa} e^1\right), \\ R &= -\left(1 - \frac{\eta_0}{\kappa}\right)^3 \Lambda'(\eta_2) e^0 \wedge e^1. \end{aligned} \quad (9)$$

For solving the field equations it is useful to use the Poisson sigma model formalism. We shall not give here the details of this formalism and refer the reader to the vast literature on the subject [9]. For our purposes it is sufficient to define a new set of variables (target space coordinates),

$$X_1 = C(\eta_A), \quad X_2 = \operatorname{arccosh} \frac{\eta_0}{\sqrt{\eta_1^2 - \eta_0^2}}, \quad X_3 = -\eta_2, \quad (10)$$

where $C(\eta_A)$ is the Casimir invariant (5) written in terms of the η_A , $C(\eta_A) = \left(1 - \frac{\eta_0}{\kappa}\right)^{-2} (\eta_1^2 - \eta_0^2) + 2 \int \Lambda(\eta_2) d\eta_2$. The transformation (10) holds when the value of the function $\gamma = C - 2 \int \Lambda(\eta_2) d\eta_2$ is negative. For positive values of γ , one must perform instead the transformation

$$X_1 = C(\eta_A), \quad X_2 = \operatorname{arcsinh} \frac{\eta_0}{\sqrt{\eta_1^2 - \eta_0^2}}, \quad X_3 = -\eta_2. \quad (11)$$

For definiteness, in the following we shall consider the case $\gamma < 0$.

The new variables satisfy the commutation relations

$$[X_1, X_2] = [X_1, X_3] = 0, \quad [X_2, X_3] = 1, \quad (12)$$

and in term of them, the η_a can be written as

$$\eta_0 = \frac{\Gamma}{\Delta} \cosh X_2, \quad \eta_1 = \frac{\Gamma}{\Delta} \sinh X_2, \quad (13)$$

where

$$\Gamma = \sqrt{\left| X_1 + 2 \int \Lambda(-X_3) dX_3 \right|}, \quad \Delta = 1 + \frac{\Gamma}{\kappa} \cosh X_2. \quad (14)$$

In these variables, the field equations take a very simple form [9]:

$$dX_1 = 0, \quad dX_2 = -B_3, \quad dX_3 = B_2, \quad dB_i = 0, \quad (15)$$

whose solutions are given by

$$X_1 = \text{const}, \quad B_1 = 2d\phi, \quad (16)$$

ϕ , X_2 and X_3 being arbitrary functions. One can then go back to the original variables. Defining $X_1 = c$, where c is the constant value of C , $X_2 = \theta$, $X_3 = -\eta$, one obtains

$$\begin{aligned} e^0 &= -\Delta \left[\frac{1}{\Gamma} \sinh \theta d\eta + \Gamma \left(\cosh \theta - \frac{\Gamma}{\kappa} \right) d\phi \right], \\ e^1 &= \Delta \left[\frac{1}{\Gamma} \cosh \theta d\eta + \Gamma \sinh \theta d\phi \right], \\ \omega &= d\theta + \Lambda d\phi \end{aligned} \quad (17)$$

where, in terms of the new variables, the functions Γ and Δ read $\Gamma = \sqrt{|c - 2 \int \Lambda d\eta|}$ and $\Delta = 1 + \Gamma \cosh \theta / \kappa$.

One is still free to choose a gauge. The most interesting choices are $\theta = 0$ or $\theta = \phi$. The first choice may be considered as the ground state of the model and yields

$$e^0 = -\Gamma \left(1 - \frac{\Gamma^2}{\kappa^2} \right) d\phi, \quad e^1 = \frac{1}{\Gamma} \left(1 + \frac{\Gamma}{\kappa} \right) d\eta, \quad \omega = \Lambda d\phi, \quad (18)$$

with

$$T^0 = \frac{3\Lambda}{\kappa} \left(1 + \frac{\Gamma}{\kappa} \right) d\phi \wedge d\eta, \quad T^1 = 0, \quad R = -\Lambda' d\phi \wedge d\eta. \quad (19)$$

In particular, for $\Lambda = \lambda\eta_2$, one obtains a deformation of (anti-)de Sitter space with constant curvature, but nonvanishing torsion.

In the second case,

$$\begin{aligned} e^0 &= -\Delta \left[\frac{1}{\Gamma} \sinh \theta d\eta + \Gamma \left(\cosh \theta - \frac{\Gamma}{\kappa} \right) d\theta \right], \\ e^1 &= \Delta \left[\frac{1}{\Gamma} \cosh \theta d\eta + \Gamma \sinh \theta d\theta \right], \\ \omega &= (1 + \Lambda) d\theta, \end{aligned} \quad (20)$$

and the components of the torsion and the curvature are

$$\begin{aligned} T^0 &= \left(-\frac{1 + \Lambda}{\kappa} \sinh^2 \theta + \frac{3}{\kappa} \Lambda \Delta \right) d\theta \wedge d\eta, \\ T^1 &= \frac{1 + \Lambda}{\kappa} \sinh \theta \cosh \theta d\theta \wedge d\eta, \\ R &= -\Lambda' d\theta \wedge d\eta. \end{aligned} \quad (21)$$

The solutions corresponding to $\gamma > 0$ can be obtained from the previous ones by simply interchanging $\sinh \theta$ with $\cosh \theta$ everywhere.

For a discussion of the properties of the solutions, it may be useful to define a line element $ds^2 = h_{ab} e^a e^b$, although this is not gauge invariant. The possibility of defining a gauge invariant metric will be discussed elsewhere [10]. Alternatively, one may interpret the lack of invariance assuming that Planck-energy particles experience different metrics depending on their state of motion. One has

$$ds^2 = \Delta^2 \left[\frac{d\eta^2}{\Gamma^2} - \Gamma^2 \left(1 - 2\frac{\Gamma}{\kappa} \cosh \theta + \frac{\Gamma^2}{\kappa^2} \right) d\theta^2 + 2\frac{\Gamma}{\kappa} \sinh \theta d\eta d\theta \right] \quad (22)$$

for $\gamma < 0$, or

$$ds^2 = \Upsilon^2 \left[\frac{-d\eta^2}{\Gamma^2} + \Gamma^2 \left(1 + 2\frac{\Gamma}{\kappa} \sinh \theta - \frac{\Gamma^2}{\kappa^2} \right) d\theta^2 + 2\frac{\Gamma}{\kappa} \cosh \theta d\eta d\theta \right] \quad (23)$$

for $\gamma > 0$, with $\Upsilon = 1 + \Gamma \sinh \theta / \kappa$.

The discussion of the properties of the solutions is in general quite complicated. First of all, we notice that the volume element is $e^0 \wedge e^1 = \Delta^3 d\eta \wedge d\theta$, which is positive definite when $\gamma < 0$. In this case, the only singularities (at least for well-behaved Λ) are horizons occurring at the zeroes of Γ . When $\gamma > 0$, instead, torsion singularities can occur at $\sinh \theta = -\kappa/\Gamma$, where Υ vanishes, and the discussion of the solutions becomes more intricate.

It can be interesting to consider as an example the case $\Lambda = \lambda\eta_2$, which corresponds to de Sitter ($\lambda < 0$) or anti-de Sitter ($\lambda > 0$) in the undeformed limit. Let us start from the case $\gamma < 0$. In this case the coordinate η is spacelike, and the solutions are time-dependent deformations of the static de Sitter or anti-de Sitter solutions, depending on the sign of λ . When λ and c have the same sign, a horizon is present at $\eta = \sqrt{c/\lambda}$, but curvature and torsion are regular everywhere.

For $\gamma > 0$, the coordinate η is timelike when $\Upsilon > 0$, and the solutions can be interpreted as space-dependent deformations of the two-dimensional cosmological de Sitter or anti-de Sitter solutions (see for example [11]). Again a horizon is present when λ and c have the same sign, but now also a singularity of the torsion occurs where $\Upsilon = 0$.

A more accurate discussion of the properties of the solutions would require the definition of a gauge-independent metric or an interpretation of its gauge dependence, which we leave for further investigations.

We have investigated a model of gravity in two dimensions, invariant under a deformation of the (anti)-de Sitter algebra or of more general algebras corresponding to dilaton-dependent potentials. We have defined the action and obtained the general solutions of the field equations. All solutions imply the presence of nontrivial torsion.

To conclude, we notice that a deformation of the (anti)-de Sitter algebra compatible with the algebra of ref. [2] in the $\lambda \rightarrow 0$ limit can be obtained also in four dimensions, using a procedure analogous to the one adopted above. The deformed algebra reads

$$\begin{aligned} [P_i, P_j] &= \lambda \left(1 - \frac{P_0}{\kappa}\right)^2 \left(\epsilon_{ijk} M_k + \frac{1}{\kappa} (N_i P_j - N_j P_i)\right), \\ [P_0, P_j] &= \lambda \left(1 - \frac{P_0}{\kappa}\right)^3 N_j, \quad [M_i, P_j] = \epsilon_{ijk} P_k, \quad [M_i, P_0] = 0, \\ [N_i, P_j] &= \delta_{ij} P_0 - \frac{P_i P_j}{\kappa}, \quad [N_i, P_0] = P_i - \frac{P_0 P_i}{\kappa}, \\ [N_i, N_j] &= \epsilon_{ijk} M_k, \quad [M_i, N_j] = \epsilon_{ijk} N_k, \quad [M_i, M_j] = \epsilon_{ijk} M_k, \end{aligned}$$

where $i, j, \dots = 1, 2, 3$, $N_i = M_{0i}$, $M_k = \frac{1}{2} \epsilon_{ijk} M_{ij}$.

Unfortunately, however, an action principle for nonlinear gauge theories is not known in four dimensions. Nevertheless, we believe that our low-dimensional model can give some hints for addressing this problem, and also other open questions, as the definition of a gauge-invariant metric and/or a suitable coupling for point particles.

References

- [1] G. Amelino-Camelia, Int. J. Mod. Phys. **D11**, 35 (2002); J. Kowalski-Glikman, Int. J. Mod. Phys. **A17**, 1 (2002); J. Lukierski, A. Nowicki, H. Ruegg and V.N. Tolstoy, Phys. Lett. **B264**, 331 (1991); S. Majid and H. Ruegg, Phys. Lett. **B334**, 348 (1994).
- [2] J. Magueijo and L. Smolin, Phys. Rev. Lett. **88**, 190403 (2002).
- [3] L. Garay, Int. J. Mod. Phys. **A10**, 145 (1995).
- [4] S. Mignemi, [hep-th/0208062](#).
- [5] S. Mignemi, Phys. Rev. **D50**, 4733 (1994), Ann. Phys. **245**, 23 (1996).
- [6] K. Isler and C. Trugenberger, Phys. Rev. Lett. **63**, 834 (1989);
- [7] D. Cangemi and R. Jackiw, Phys. Rev. Lett. **69**, 233 (1992).
- [8] N. Ikeda, Ann. Phys. **235**, 435 (1994).
- [9] P. Schaller and T. Strobl, Mod. Phys. Lett. **A9**, 3129 (1994); T. Klösch and T. Strobl, Class. Quantum Grav. **13**, 965 (1996); D. Grumiller, W. Kummer and D.V. Vassilevich, [hep-th/0204253](#).
- [10] D. Grumiller, W. Kummer, S. Mignemi, T. Strobl and D.V. Vassilevich, in preparation.
- [11] M. Cadoni and S. Mignemi, Gen. Relativ. Grav. **34**, in press.